

Padena Fruit Set

Most of summer fruits needs heavily good stored nutrients to kick-start next year's growth in the spring. Nitrogen, Potassium, Zinc and Boron along with seaweed extract are recommended for this application. Seaweed content of Padena Fruit Set, enhance the abilities of absorption, translocation and metabolisms of nutritional elements in the plant. This is achieved through a range of beneficial properties as associated with this bio-stimulant which interact directly with the biologic system of the plant.

Supplying the adequate amounts of each element, make an opportunity for a good and strong start on next season. For this, Padena Fertilizer Co., introduce PADENA FRUIT SET as a creative and smart choice on Post Harvesting and early growing stages.

Benefits

- Rapid absorption for better and faster penetration and translocation of elements in leaves and phloem.
- Activation of vital enzymatic systems according to short remaining time of plant activity.
- Rapid production and translocation of polysaccharides in plants.
- Induce the protein production and vitality of buds for better hibernation.
- Increasing the blooming and fruit production on next season.
- Inducing the tolerance to early season stresses, including cold weather shocks.
- Increases resistance to cold of winter and early spring
- Miscible with most pesticides and fertilizers

Composition (W/V)%

Zn	6
B	0.5
N	3

K₂O	6
Seaweed extract	4

Application Rate

Crop	How to use	Dosage	Time
Seed trees	Foliar Application	2-3 liters per 1000 liters of water per hectare	14 days after harvest and repeat at the time of swelling of next year's reproductive buds
Fruit trees	Foliar Application	2-3 liters per 1000 liters of water per hectare	Mid-autumn and repeat at the time of swelling of reproductive buds next year
Pistachio	Foliar Application	5 liters per 1000 liters of water per hectare	14 days after harvest and repeat at the time of swelling of next year's reproductive buds
Pomegranate	Foliar Application	2 liters per 1000 liters of water per hectare	14 days after harvest and repeat at the time of swelling of next year's reproductive buds
Kiwi fruit	Foliar Application	3 liters per 1000 liters of water per hectare	After harvest and repeat at the time of swelling of next year's reproductive buds